

ON A RECURSIVE EQUATION OVER p -ADIC FIELD

FARRUKH MUKHAMEDOV

ABSTRACT. In the paper we completely describe the set of all solutions of a recursive equation, arising from the Bethe lattice models over p -adic numbers.

Mathematics Subject Classification: 46S10, 12J12.

Key words: recursive equation, p -adic numbers, model, fixed point, unique.

1. INTRODUCTION

The p -adic numbers were introduced by K. Hensel. For about a century after the inventing of the p -adic numbers, they were mainly considered objects of pure mathematics. After discovering that the physics of certain models could be based on the idea that the structure of space-time for very short distances might conveniently be described in terms of p -adic numbers, many applications of such numbers in theoretical physics have been proposed in works (see for example, [4],[6],[12],[15]). A number of p -adic models in physics cannot be described using ordinary probability theory based on the Kolmogorov axioms, therefore p -adic probability models were investigated in [7]. Such models appear to provide the probabilistic interpretation of p -adic valued wave functions and string amplitudes in the framework of p -adic theoretical physics (see [6, 7]). Using such a p -adic probability approach in [13, 14] we have developed a theory of statistical mechanics in the context of that theory of probability. In statistical mechanics, due to their solvable character [1], Bethe lattice models and models on random tree-like graphs [3], play a central role. In the calculation of the average magnetization of these models naturally appears the following recursive equation

$$h_n = \log \left[\left(\frac{e^{\alpha_{n+1} + h_{n+1}} + e^{\beta_{n+1}}}{e^{\gamma_{n+1}} + e^{h_{n+1}}} \right) \left(\frac{e^{\alpha_{n+2} + h_{n+2}} + e^{\beta_{n+2}}}{e^{\gamma_{n+2}} + e^{h_{n+2}}} \right) \right], \quad n \in \mathbb{N}, \quad (1.1)$$

here $\alpha_k, \beta_k, \gamma_k \in \mathbb{R}$ and $\{h_n\}$ is an unknown sequence of real numbers, the solution of such a equation describes corresponding Gibbs measures. On certain conditions depending on parameters $\alpha_k, \beta_k, \gamma_k$ the equation (1.1) has infinitely many solutions [2],[5] (i.e. over real numbers), but all the set of solutions is still not described.

On leave from Department of Mechanics and Mathematics, National University of Uzbekistan, Tashkent, 700174, Uzbekistan.

In the present paper we are going to completely describe the set of solutions of (1.1) over p -adic numbers. Our hope is that this allows us further understanding of the Bethe lattice models and models on random tree-like graphs over p -adic numbers. Moreover, we think that the obtained result will give certain information on random rational p -adic dynamical systems generated by linear-fractional functions. Note that some applications of p -adic dynamical systems to some biological and physical systems have been proposed in [8].

Throughout the paper p will be a fixed prime number greater than 3, i.e. $p \geq 3$. Every rational number $x \neq 0$ can be represented in the form $x = p^r \frac{n}{m}$, where $r, n \in \mathbb{Z}$, m is a positive integer, $(p, n) = 1$, $(p, m) = 1$. The p -adic norm of x is given by

$$|x|_p = \begin{cases} p^{-r} & \text{for } x \neq 0 \\ 0 & \text{for } x = 0. \end{cases}$$

It satisfies the following strong triangle inequality

$$|x + y|_p \leq \max\{|x|_p, |y|_p\},$$

this is a non-Archimedean norm.

The completion of the field of rational numbers $\mathbb{Q}$ with respect to the p -adic norm is called *p -adic field* and it is denoted by $\mathbb{Q}_p$.

Given $a \in \mathbb{Q}_p$ and $r > 0$ put

$$B(a, r) = \{x \in \mathbb{Q}_p : |x - a|_p < r\}, \quad S(a, r) = \{x \in \mathbb{Q}_p : |x - a|_p = r\}.$$

The *p -adic logarithm* is defined by the series

$$\log_p(x) = \log_p(1 + (x - 1)) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(x - 1)^n}{n},$$

which converges for $x \in B(1, 1)$; the *p -adic exponential* is defined by

$$\exp_p(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!},$$

which converges for $x \in B(0, p^{-1/(p-1)})$.

Lemma 1.1. [9] *Let $x \in B(0, p^{-1/(p-1)})$ then we have*

$$|\exp_p(x)|_p = 1, \quad |\exp_p(x) - 1|_p = |x|_p, \quad |\log_p(1 + x)|_p = |x|_p \quad (1.2)$$

$$\log_p(\exp_p(x)) = x, \quad \exp_p(\log_p(1 + x)) = 1 + x. \quad (1.3)$$

2. MAIN RESULT

In this section we are going to completely describe the set of solutions of (1.1) over p -adic numbers. Let us first formulate the problem in a p -adic setting. Namely,

we are interested to find all solutions of the following recursive equation

$$h_n = \log_p \left[\left(\frac{\exp_p(\alpha_{n+1} + h_{n+1}) + \exp_p(\beta_{n+1})}{\exp_p(\gamma_{n+1}) + \exp_p(h_{n+1})} \right) \left(\frac{\exp_p(\alpha_{n+2} + h_{n+2}) + \exp_p(\beta_{n+2})}{\exp_p(\gamma_{n+2}) + \exp_p(h_{n+2})} \right) \right], \quad n \in \mathbb{N}, \quad (2.1)$$

here $\alpha_k, \beta_k, \gamma_k \in \mathbb{Q}_p$, such that

$$|\alpha_k|_p < p^{-1/(p-1)}, \quad |\beta_k|_p < p^{-1/(p-1)}, \quad |\gamma_k|_p < p^{-1/(p-1)}, \quad (2.2)$$

and $h_k \in \mathbb{Q}_p$ also should satisfy $|h_k|_p < p^{-1/(p-1)}$ for every $k \in \mathbb{N}$, since these conditions ensure the existence of the p -adic $\log_p$ and $\exp_p$.

Denote

$$S = \{\mathbf{h} = (h_n)_{n \in \mathbb{N}} : h_n \text{ satisfies (2.1)}\}.$$

Let us rewrite (2.1) as follows

$$u_n = \left(\frac{a_{n+1}u_{n+1} + b_{n+1}}{c_{n+1} + u_{n+1}} \right) \left(\frac{a_{n+2}u_{n+2} + b_{n+2}}{c_{n+2} + u_{n+2}} \right), \quad n \in \mathbb{N}, \quad (2.3)$$

here we have denoted

$$\left. \begin{aligned} a_k &= \exp_p(\alpha_k), & b_k &= \exp_p(\beta_k), \\ c_k &= \exp_p(\gamma_k), & u_k &= \exp_p(h_k). \end{aligned} \right\}, \quad k \in \mathbb{N}. \quad (2.4)$$

Let us introduce

$$f_k(x) = \frac{a_k x + b_k}{c_k + x}, \quad k \in \mathbb{N}. \quad (2.5)$$

The functions f_k have the the following properties.

Lemma 2.1. *For every $k \in \mathbb{N}$ the following relations hold*

$$|f_k(x) - f_k(y)|_p \leq \frac{1}{p} \frac{|x - y|_p}{|c_k + x|_p |c_k + y|_p} \quad (2.6)$$

$$|f_k(\exp_p(h))|_p = 1 \quad \text{for every } |h|_p \leq \frac{1}{p} \quad (2.7)$$

Proof. From (2.5) we immediately obtain

$$|f_k(x) - f_k(y)|_p = \frac{|x - y|_p |a_k c_k - b_k|_p}{|c_k + x|_p |c_k + y|_p}.$$

Keeping in mind (2.4) from (1.2),(2.2) one gets

$$|a_k c_k - b_k|_p = |\exp_p(\alpha_k + \gamma_k - \beta_k) - 1|_p \leq \frac{1}{p}$$

which implies (2.6). The strong triangle inequality with (1.2) and $p \geq 3$ implies that $|\exp_p(h) + 1|_p = 1$, hence one finds (2.7). $\square$

The main result of the paper is the following

Theorem 2.2. *Let (2.2) be satisfied. Then $|S| \leq 1$, here $|A|$ means the number of elements of a set A .*

Proof. If $S = \emptyset$, then nothing to prove. So, assume that $S \neq \emptyset$. To prove Theorem it is enough to prove that any two elements of S coincide with each other. To show this it is enough to prove that for arbitrary $\varepsilon > 0$ and every $\mathbf{h} = (h_n, n \in \mathbb{N})$, $\mathbf{s} = (s_n, n \in \mathbb{N}) \in S$ and $n \in \mathbb{N}$ the inequality $|h_n - s_n|_p < \varepsilon$ is valid.

Let $\mathbf{h} = (h_n, n \in \mathbb{N})$, $\mathbf{s} = (s_n, n \in \mathbb{N}) \in S$ and $\varepsilon > 0$ be an arbitrary number. Put $v_k = \exp_p(s_k)$. Let $n \in \mathbb{N}$ be an arbitrary number. Using (2.5), (2.3) and Lemma 2.1 we get

$$\begin{aligned} |u_n - v_n|_p &= |f_{n+1}(u_{n+1})f_{n+2}(u_{n+2}) - f_{n+1}(v_{n+1})f_{n+2}(v_{n+2})|_p \\ &\leq \max \left\{ |f_{n+1}(u_{n+1})|_p |f_{n+2}(u_{n+2}) - f_{n+2}(v_{n+2})|_p, \right. \\ &\quad \left. |f_{n+2}(v_{n+2})|_p |f_{n+1}(u_{n+1}) - f_{n+1}(v_{n+1})|_p \right\} \\ &\leq \frac{1}{p} \max\{|u_{n+1} - v_{n+1}|_p, |u_{n+2} - v_{n+2}|_p\}, \end{aligned} \quad (2.8)$$

here we have used the following equalities

$$|c_k + u_k|_p = 1, \quad |c_k + v_k|_p = 1, \quad \forall k \in \mathbb{N},$$

which immediately follow from (1.2).

Now take $n_0 \in \mathbb{N}$ such that $\frac{1}{p^{n_0}} < \varepsilon$. Then iterating (2.8) one gets

$$|u_n - v_n|_p \leq \frac{1}{p^{n_0}} < \varepsilon$$

The last inequality with (1.2), (2.4) implies that

$$|h_n - s_n|_p = |u_n - v_n|_p < \varepsilon$$

This completes the proof. $\square$

Remark. Note that the proved Theorem provides some applications to the p -adic Bethe lattice models and models on random tree-like graphs. This would be a theme of our further investigations. We also think that the result can be applied for the study of dynamics of random rational p -adic dynamical systems. It should be also noted that certain type of recursive equations in p -adic numbers were considered in [11], they were related with transcendentality of the p -adic numbers. On the other hand, the method presented here used functional approach which is different from [11], where some number theoretical method was presented.

Now consider several cases when S is non empty.

Case 1. Let us assume that the following equality holds

$$a_k + b_k = c_k + 1$$

for every $k \in \mathbb{N}$, which implies that the equation (2.3) has a solution $u_n = 1$ for all $n \in \mathbb{N}$. According to Theorem 2.2 it is unique, i.e. $|S| = 1$.

Case 2. Now suppose that $a_k = a$, $b_k = b$, $c_k = c$ for all $k \in \mathbb{N}$. We are going to show the existence of a solution (2.3). Let us search a solution in a form $u_n = u$, $\forall n \in \mathbb{N}$, here u is unknown. Then (2.3) reduces to the equation $f(u) = u$, where

$$f(u) = \left(\frac{au + b}{c + u} \right)^2.$$

Put $D = S(0, 1) \cap B(1, p^{-1/(p-1)})$. Let us show that $f(D) \subset D$. Indeed, let $x \in D$, then $|x|_p = 1$, $|x - 1|_p < p^{-1/(p-1)}$. According to Lemma 1.1 this means that $x = \exp_p(y)$ for some $y \in B(0, p^{-1/(p-1)})$. Now by means of Lemma 2.1 we infer that $|f(x)|_p = 1$. The strong triangle inequality with

$$|a - 1|_p \leq \frac{1}{p}, \quad |b - c| \leq \frac{1}{p}$$

implies that

$$|f(x) - 1|_p = \left| \frac{(a-1)x + b - c}{c + x} \right|_p \leq \frac{1}{p}. \quad (2.9)$$

Using the same argument along the proof of Lemma 2.1 one gets

$$|f(x) - f(y)|_p \leq \frac{1}{p} |x - y|_p \quad \text{for every } x, y \in D. \quad (2.10)$$

Thus from the inequalities (2.9), (2.10) we obtain that f is a contraction of D , hence f has a unique fixed point $\zeta \in D$. Consequently, S is non empty. Now Theorem 2.2 yields that $|S| = 1$.

Acknowledgement. The author thanks the FCT (Portugal) grant SFRH/BPD/17419/2004, and also express gratitude to referees for useful suggestions.

REFERENCES

- [1] R.J. Baxter, *Exactly Solved Models in Statistical Mechanics*, (Academic Press, London/New York, 1982).
- [2] P.M. Bleher, N.N.Ganikhodjaev, Pure phases of the Ising model on Bethe lattices. *Theory Probab. Appl.* **35** (1991), 2, 216–227.
- [3] S.N. Dorogovtsev, J.F.F. Mendes, *Evolution of networks: from biological nets to the internet and WWW*. (Oxford Univ. Press 2003).
- [4] P.G.O.Freund, E.Witten, Adelic string amplitudes, *Phys. Lett.* **B199**, 191-194 (1987).
- [5] H.O. Georgii, *Gibbs measures and phase transitions* (Walter de Gruyter, Berlin, 1988).
- [6] A.Yu.Khrennikov, *p-adic Valued Distributions in Mathematical Physics*. (Kluwer Academic Publisher, Dordrecht, 1994).
- [7] A.Yu.Khrennikov, *Non-Archimedean analysis: quantum paradoxes, dynamical systems and biological models*. (Kluwer Academic Publisher, Dordrecht, 1997).
- [8] A.Yu. Khrennikov, M. Nilsson M., *p-adic deterministic and random dynamical systems*. (Kluwer, Dordrecht, 2004).

- [9] N.Koblitz, *p-adic numbers, p-adic analysis and zeta-function*, Berlin, Springer, 1977.
- [10] R. Lyons, The Ising model and percolation on trees and tree-like graphs, *Commun. Math. Phys.* **125**(1989), 337–353.
- [11] K. Mahler, *p-adic numbers and their functions*. Cambridge Tracts in Mathematics, **76**, (Cambridge Univ. Press, Cambridge-New York, 1981).
- [12] E.Marinary, G.Parisi, On the p -adic five point function, *Phys. Lett.* **203B** 52-56 (1988).
- [13] F.M.Mukhamedov, U.A.Rozikov, On Gibbs measures of p -adic Potts model on the Cayley tree, *Indag. Math. N.S.* **15**(2004), 85–100.
- [14] F.M.Mukhamedov, U.A.Rozikov, On inhomogeneous p -adic Potts model on a Cayley tree, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **8** (2005) 277–290.
- [15] V.S.Vladimirov, I.V.Volovich, E.I.Zelenov, *p-adic Analysis and Mathematical Physics*, (World Scientific, Singapour, 1994).

FARRUKH MUKHAMEDOV, DEPARTAMENTO DE FISICA, UNIVERSIDADE DE AVEIRO,
 CAMPUS UNIVERSITÁRIO DE SANTIAGO, 3810-193 AVEIRO, PORTUGAL
E-mail address: far75m@yandex.ru farruh@fis.ua.pt